

how to inject subcutaneously

what you'll need

Reconstituted peptide · Insulin syringe · Alcohol swabs · Sharps bin

- 1** **prepare your dose**
Draw your prescribed dose from the peptide vial using a clean insulin syringe. Remove air bubbles by gently tapping and pushing the plunger up slowly.
- 2** **choose your injection site**
Best sites: lower abdomen (2-3 cm from navel), outer thigh, or back of upper arm. Rotate sites with each injection.
- 3** **clean the site**
Wipe the chosen area with an alcohol swab. Let it air dry for 10 seconds - do not blow on it or wipe it off.
- 4** **pinch and insert**
Gently pinch 2-3 cm of skin. Insert the needle at a 45° angle with a smooth, steady motion. Release the pinch once the needle is in.
- 5** **inject slowly**
Push the plunger down slowly and steadily. Minimal resistance is normal. If you feel strong pain or burning, stop and reposition.
- 6** **remove and dispose**
Pull the needle out smoothly at the same angle. Place immediately into a sharps bin - never recap. Apply gentle pressure with a clean swab. Do not rub.

best injection sites

Lower abdomen · Outer thigh · Back of upper arm · Always rotate

important

Never inject into a vein, muscle, or broken skin. If you notice redness, swelling, or unusual pain, stop and contact us.



how to reconstitute your peptide

what you'll need

Peptide vial · Bacteriostatic water · 1 ml insulin syringe · Alcohol swabs

- 1** wash your hands
Wash thoroughly with soap and water before touching anything.
Cleanliness is the most important part of this process.
- 2** wipe both vial tops with an alcohol swab
Swab the rubber stopper on your peptide vial and BAC water vial.
Let air dry for 10-15 seconds - do not blow on them.
- 3** draw BAC water into the syringe
Push the needle into the BAC water vial and slowly pull the plunger to draw the required amount. 1-2 ml is standard for most peptides.
- 4** inject water slowly into the peptide vial
Aim the water at the glass wall - not directly onto the powder.
Push the plunger very slowly and gently.
- 5** let it dissolve
Remove the syringe and let the vial sit. Gently roll between your palms if needed. Never shake it.

important

Always inject water gently against the glass wall - never directly onto the powder. Peptides are delicate; aggressive mixing can damage them.

